4. Other Discretization Methods

4.1. Discontinuous Galerkin (DG) Methods = Non-conforming FEM

4.1.1. Derivation of the DG Variational Formulation and the DG Schemes

- Let us again consider the model problem

(1) Given $f \in L_2(\Omega)$, find $u \in \bar{V} = H^1(\Omega)$ such that

$$-\Delta u = f \quad \text{in } \Omega \subset \mathbb{R}^d \quad (\Gamma \wedge \text{Lip}, d = 2,3)$$
$$u = g \quad \text{on } \Gamma = \partial\Omega$$

- Solvability:

Due to Lax-Milgram, there exists a unique weak solution $u \in \bar{V}_0 = \mathring{H}^1(\Omega) \subset V = H^1(\Omega)$:

$$(1)_{VF} \quad \int_\Omega \nabla u \cdot \nabla v \, dx = \int_\Omega f v \, dx \quad \forall v \in \mathring{H}^1(\Omega)$$
$$\| \qquad\qquad \| \qquad\qquad \| =: \langle F, v \rangle \quad \forall v \in \bar{V}_0$$
$$a(u,v) \qquad = (f,v)_{0,\Omega}$$

- Exercise 4.1: (see also Chapter 2)

Show that $\nabla u \in H(\text{div}, \Omega)$ and $\text{div}\, \nabla u = -f \in L_2(\Omega)$!

Therefore, $u \in \mathring{H}^1(\Omega) \wedge H^\Delta(\Omega)$, where

$$H^\Delta(\Omega) := \{ u \in H^1(\Omega) : \Delta u = \text{div}\, \nabla u \in L_2(\Omega) \}$$
$$= \{ u \in H^1(\Omega) : \nabla u \in H(\text{div}, \Omega) \}.$$