

# T U T O R I A L

## “Numerical Methods for the Solution of Elliptic Partial Differential Equations”

to the lecture

“Numerical Methods for Elliptic Problems”

**Tutorial 7**    Tuesday, 12 May 2009 (Time : 10<sup>15</sup> – 11<sup>45</sup>    Room : T 911 )

### 2.5 Properties of the Finite Elements Equations

**31** Prove the inheritance identity

$$(K_h \underline{u}_h, \underline{v}_h) = a(u_h, v_h) \quad \forall \underline{u}_h, \underline{v}_h \leftrightarrow u_h, v_h \in V_{0h} ! \quad (2.29)$$

**32** Show that the eigenvalue estimates in Theorem 2.4 are sharp with respect to the  $h$ -order by proving the following statement. There exist positive constants  $\underline{c}'_E$  and  $\bar{c}'_E$  independent of  $h$  satisfying the estimates

$$\lambda_{\min}(K_h) \leq \underline{c}'_E h^d \quad \text{and} \quad \lambda_{\max}(K_h) \geq \bar{c}'_E h^{d-2}. \quad (2.30)$$

For simplicity, consider the 1D case ( $d = 1$ ):

$$\begin{aligned} -u''(x) &= f(x) & \forall x \in (0, 1), \\ u(0) &= u(1) = 0. \end{aligned}$$

**33** Show that, for a regular triangulation according to Definition 2.3, there exist  $h$ -independent positive constants  $\underline{c}_0$  and  $\bar{c}_0$  satisfying the inequalities

$$\underline{c}_0 h^d (\underline{v}_h, \underline{v}_h) \leq (M_h \underline{v}_h, \underline{v}_h) \leq \bar{c}_0 h^d (\underline{v}_h, \underline{v}_h) \quad (2.31)$$

for all  $\underline{v}_h \in \mathbb{R}^{N_h}$ , where  $M_h$  denotes the mass-matrix defined by the identity

$$(M_h \underline{u}_h, \underline{v}_h) := \int_{\Omega} u_h(x) v_h(x) dx \quad \forall \underline{u}_h, \underline{v}_h \leftrightarrow u_h, v_h \in V_{0h}. \quad (2.32)$$

**34** Let  $\lambda = \lambda_{\max}$  be the maximal eigenvalue of the generalized eigenvalue problem

$$K_h \underline{u}_h = \lambda M_h \underline{u}_h \quad (2.33)$$

and let  $\lambda_r = \lambda_{r, \max}$  be the maximal eigenvalues of generalized eigenvalue problems

$$K_h^{(r)} \underline{u}_h^{(r)} = \lambda_r M_h^{(r)} \underline{u}_h^{(r)}, \quad (2.34)$$

where  $K_h^{(r)}$  and  $M_h^{(r)}$  denote the (local) element stiffness and mass matrices for element number  $r = 1, 2, \dots, R_h$ , i. e., it holds

$$K_h = \sum_{r=1}^{R_h} C_r K_h^{(r)} C_r^T \quad \text{and} \quad M_h = \sum_{r=1}^{R_h} C_r M_h^{(r)} C_r^T .$$

Show the eigenvalue estimate

$$\lambda \leq \max_{r=1,2,\dots,R_h} \lambda_r . \tag{2.35}$$