<u>TUTORIAL</u>

"Computational Mechanics"

to the lecture

"Numerical Methods in Continuum Mechanics 1"

Tutorial 01 Friday, March 12, 2010 (Time : $10^{15} - 11^{00}$ Room : HS 12)

1 Introduction to Mixed Variational Formulations: Examples

1.1 Scalar Elliptic BVP of Second Order

01 Provide the mixed variational formulation of the mixed BVP

$$-\Delta u = f \text{ in } \Omega, \ u = g_1 \text{ on } \Gamma_1, \ \frac{\partial u}{\partial n} = g_2 \text{ on } \Gamma_2$$

for the Poisson equation with given f, g_1, g_2, Γ_1 and Γ_2 , where $\Gamma_1 \cap \Gamma_2 = \emptyset$ and $\Gamma_1 \cup \Gamma_2 = \Gamma = \partial \Omega$!

1.2 The Stokes Equations

02 Let us consider a two-dimensional, steady state (stationary) flow of a highly viscous, incompressible fluid that can be described by the Stokes Equations

$$-\nu\Delta u + \nabla p = f \quad \text{in } \Omega, \tag{1.1}$$

$$\operatorname{div} u = 0 \quad \text{in } \Omega. \tag{1.2}$$

Assume that the velocity u can be represented by a so-called (scalar) stream function ψ as

$$u = \operatorname{curl} \psi \tag{1.3}$$

with

$$\mathbf{curl}\,\psi = \begin{pmatrix} \frac{\partial\psi}{\partial x_2} \\ -\frac{\partial\psi}{\partial x_1} \end{pmatrix}$$

Show that

$$\operatorname{div} u = 0, \tag{1.4}$$

and derive the following relation from the equations (1.1):

$$\nu \Delta^2 \psi = \operatorname{curl} f \quad \text{in } \Omega, \tag{1.5}$$

where the so-called scalar curl is now defined by the relation

$$\operatorname{curl} f = \frac{\partial f_2}{\partial x_1} - \frac{\partial f_1}{\partial x_2}.$$
(1.6)

<u>Hint:</u> See Example 1.3 from the lecture for possible boundary conditions which can be prescribed for the biharmonic equation !