

01 Show that every linear second order partial differential equation

$$-(a(x) u'(x))' + b(x) u'(x) + c(x) u(x) = f(x),$$

with $a \in \mathcal{C}^1(0, 1)$ and $b, c \in \mathcal{C}(0, 1)$ can be rewritten in the form

$$\bar{a}(x) u''(x) + \bar{b}(x) u'(x) + c(x) u(x) = f(x),$$

and find suitable functions $\bar{a} \in \mathcal{C}^1(0, 1)$ and $\bar{b} \in \mathcal{C}(0, 1)$. Show also the reverse direction.

02 Derive the variational formulations of the two following boundary value problems:

$$\begin{aligned} \text{(a)} \quad & \begin{cases} -u''(x) + u(x) = f(x) & \text{for } x \in (0, 1) \\ u(0) = g_0 \\ u(1) = g_1 \end{cases} \\ \text{(b)} \quad & \begin{cases} -u''(x) + u(x) = f(x) & \text{for } x \in (0, 1) \\ u(0) = g_0 \\ u'(1) = g_1 - \alpha_1 u(1) \end{cases} \end{aligned}$$

In particular, specify the spaces V_g , and V_0 , the bilinear form $a(\cdot, \cdot)$, and the linear form $\langle F, \cdot \rangle$.

Hint for (b): Perform integration by parts as usual, substitute $u'(1)$ due to the Robin boundary condition, and collect the bilinear and linear terms accordingly.

03 Consider the boundary value problem

$$\begin{aligned} -(a(x) u'(x))' &= 1 & \text{for } x \in (0, 1), \\ u(0) &= 0, \\ a(1) u'(1) &= 0, \end{aligned} \tag{1.1}$$

where $a(x) = \sqrt{2x - x^2}$. Justify that $u(x) = \sqrt{2x - x^2}$ is a *classical* solution of (1.1), i. e., $u \in X := \mathcal{C}^2(0, 1) \cap \mathcal{C}^1(0, 1] \cap \mathcal{C}[0, 1]$. Furthermore, show that

$$\int_0^1 |u'(x)|^2 dx = \infty.$$

Note: This example shows that $u \notin H^1(0, 1)$, i. e., u is no *weak* solution.

04 Consider the piecewise constant coefficient function $a \in L^\infty(0, 1)$,

$$a(x) = \begin{cases} a_1 & \text{for } x \in [0, \frac{1}{2}], \\ a_2 & \text{for } x \in (\frac{1}{2}, 1], \end{cases}$$

with positive constants $a_1 \neq a_2$. Derive a variational formulation for the boundary value problem

$$\begin{aligned} -a(x) u''(x) &= f(x) & \text{for } x \in (0, 1) \setminus \{\frac{1}{2}\}, \\ u(0) &= g_1, & u(1) = g_2, \end{aligned}$$

together with the *transmission conditions*

$$u(\frac{1}{2}^-) = u(\frac{1}{2}^+), \quad a_1 u'(\frac{1}{2}^-) = a_2 u'(\frac{1}{2}^+),$$

where $w(\frac{1}{2}^-)$ and $w(\frac{1}{2}^+)$ denote the left sided and right sided limit of a function w , respectively.

Hint: Integration by parts is only valid on subintervals!

05 Let the sequence $(u_k)_{k \in \mathbb{N}}$ of functions be defined by

$$u_k(x) = \begin{cases} 2x & \text{for } x \in [0, \frac{1}{2} - \frac{1}{2k}], \\ 1 - \frac{1}{2k} - 2k(x - \frac{1}{2})^2 & \text{for } x \in (\frac{1}{2} - \frac{1}{2k}, \frac{1}{2} + \frac{1}{2k}), \\ 2(1-x) & \text{for } x \in [\frac{1}{2} + \frac{1}{2k}, 1]. \end{cases}$$

Show that $u_k \in C^1[0, 1]$. Let u be defined by

$$u(x) = \begin{cases} 2x & \text{for } x \in [0, \frac{1}{2}], \\ 2(1-x) & \text{for } x \in (\frac{1}{2}, 1]. \end{cases}$$

Find out if $u, u_k \in H^1(0, 1)$ or not and justify your answer. Calculate $\|u_k - u\|_{H^1(0,1)}$ (maybe with a little help from Mathematica/Maple) or find a suitable bound for it in order to show that

$$\lim_{k \rightarrow \infty} \|u_k - u\|_{H^1(0,1)} = 0.$$

Use these results to show that $(u_k)_{k \in \mathbb{N}}$ is a Cauchy sequence in $C^1[0, 1]$ with respect to the H^1 -norm, but that there exists no limit in $C^1[0, 1]$.

06 Show that there exists **no** function $w \in L^2(0, 1)$ such that

$$\varphi(\frac{1}{2}) = \int_0^1 w(x) \varphi(x) dx \quad \text{for all } \varphi \in C_0^\infty(0, 1).$$

Hint: Consider the sequence of test functions

$$\varphi_n(x) := \begin{cases} e^{1 - \frac{1}{1-n^2(1-2x)^2}} & \text{for } |1-2x| < \frac{1}{n}, \\ 0 & \text{else} \end{cases} \in C_0^\infty(0, 1) \quad \text{for } n \in \mathbb{N}, n \geq 2.$$