

# T U T O R I A L

## “Numerical Methods for the Solution of Elliptic Partial Differential Equations”

to the lecture

“Numerics of Elliptic Problems”

### **Tutorial 05**

Tuesday, 24 April 2018, Time: 10<sup>15</sup> – 11<sup>45</sup>, Room: S2 346.

## 3 Galerkin FEM

### 3.1 Galerkin-Ritz-Method

**23** Let us consider the variational problem: Find  $u \in V_g = V_0 = L_2(0, 1)$ :

$$\int_0^1 u(x)v(x) dx = \int_0^1 f(x)v(x) dx \quad \forall v \in V_0. \quad (3.15)$$

Solve this variational problem with the Galerkin-Method using the basis

$$V_{0h} = V_{0n} = \text{span}\{1, x, x^2, \dots, x^{n-1}\},$$

where the right-hand side is given as  $f(x) = \cos(k\pi x)$ ,  $k = l + 1$  and  $l$  is the last digit from your study code (Matrikelnummer) ! Compute the stiffness matrix  $K_h$  analytically and solve the linear system  $K_h \underline{u}_h = \underline{f}_h$  numerically using the Gaussian elimination method ! Consider  $n$  to be 2, 4, 8, 10, 50, 100 !

### 3.2 Generation of the System of Finite Element Equations

**24** Show that the integration rule

$$\int_{\Delta} f(\xi, \eta) d\xi d\eta \approx \frac{1}{2} \{ \alpha_1 f(\xi_1, \eta_1) + \alpha_2 f(\xi_2, \eta_2) + \alpha_3 f(\xi_3, \eta_3) \} \quad (3.16)$$

integrates quadratic polynomials exactly, if the the weights  $\alpha_i$  and the integration points  $(\xi_i, \eta_i)$  are chosen as follows:  $\alpha_1 = \alpha_2 = \alpha_3 = 1/3$  und  $(\xi_1, \eta_1) = (1/2, 0)$ ,  $(\xi_2, \eta_2) = (1/2, 1/2)$ ,  $(\xi_3, \eta_3) = (0, 1/2)$ .

**Hint:** cf. also Exercise 17 !

**25** Let us assume that  $\mathcal{T}_h = \{\delta_r : r \in \mathbb{R}_h\}$  is a regular triangulation of the polygonally bounded Lipschitz domain  $\bar{\Omega} = \cup_{r \in \mathbb{R}_h} \bar{\delta}_r \subset \mathbb{R}^2$  into triangles  $\delta_r$ , and let  $u \in H^2(\Omega)$ . Let us now compute the integral

$$I(u) = \int_{\Omega} u(x) dx$$

by the quadrature rule

$$I_h(u) = \sum_{r \in \mathbb{R}_h} u(x_{\delta_r}(\xi^*)) |\delta_r|,$$

where  $x_{\delta_r}(\cdot)$  maps the unit triangle  $\Delta$  onto  $\delta_r$ , and  $\xi^* = (1/3, 1/3)$ . Show that

$$|I(u) - I_h(u)| \leq ch^2 |u|_{H^2(\Omega)},$$

where  $c$  is some generic positive constant. Can you weaken the assumption that  $u \in H^2(\Omega)$  ?

**Hint:** Use the mapping principle and the Bramble-Hilbert Lemma; cf. also Exercise 17 !

26 Show the inequality

$$\frac{1}{2} \sin \theta_r h_r^2 \leq |J_{\delta_r}| \leq \frac{\sqrt{3}}{2} h_r^2, \quad (3.17)$$

where  $h_r$  is the largest edge and  $\theta_r$  the smallest angle of the triangle  $\delta_r$ .