

Die Koeffizienten von

$$K^{(i)} = \begin{bmatrix} K_{11}^{(i)} & K_{12}^{(i)} & K_{13}^{(i)} \\ K_{21}^{(i)} & K_{22}^{(i)} & K_{23}^{(i)} \\ K_{31}^{(i)} & K_{32}^{(i)} & K_{33}^{(i)} \end{bmatrix} \quad \text{und} \quad \underline{f}^{(i)} = \begin{bmatrix} f_1^{(i)} \\ f_2^{(i)} \\ f_3^{(i)} \end{bmatrix}$$

werden mittels Abbildung auf das Basiselement

$$\bar{\delta}_i = [x_{i-1}, x_i] \xrightleftharpoons[\alpha = x_{\delta_i(\eta)}]{\eta = \xi \delta_i(x)} \bar{\Delta} = [0, 1]$$

berechnet, d.h. für unser Bsp (vgl. (2)) $\lambda(x) = 1$

$$\int_a^b \lambda(x) u_h'(x) v_h'(x) dx = \sum_{i=1}^n \int_{x_{i-1}}^{x_i} \lambda(x) u_h'(x) v_h'(x) dx$$

folgt z.B. für

$$K_{12}^{(i)} = \int_{x_{i-1}}^{x_i} \lambda(x) \frac{d\varphi_{2i-2}(x)}{dx} \frac{d\varphi_{2i-2}(x)}{dx} dx$$

$$= \int_0^1 \lambda(x_{\delta_i(\eta)}) \frac{1}{h} \frac{d\phi_1(\eta)}{d\eta} \frac{1}{h} \frac{d\phi_1(\eta)}{d\eta} h d\eta$$

$$\delta_i \leftrightarrow \Delta : x = (x_i - x_{i-1}) \eta + x_{i-1} = h\eta + x_{i-1}$$

$$\eta = \frac{x - x_{i-1}}{h}$$

$$dx = h d\eta \quad \frac{dx}{d\eta} = \frac{1}{\frac{d\eta}{dx}} \frac{d\eta}{dx} = \frac{1}{h} \frac{d\eta}{dx}$$

$$= \frac{1}{h} \int_0^1 \underbrace{\lambda(x_{\delta_i(\eta)}) \frac{d\phi_1(\eta)}{d\eta} \frac{d\phi_1(\eta)}{d\eta}}_{=: q_i(\eta)} d\eta$$

$$\approx \frac{1}{h} \left[\frac{1}{2} q_i\left(\frac{1}{2} \left(1 - \frac{1}{\sqrt{3}}\right)\right) + \frac{1}{2} q_i\left(\frac{1}{2} \left(1 + \frac{1}{\sqrt{3}}\right)\right) \right]$$

GAUSS 2: