## Numerical Methods for Partial Differential EquationsWS 2010 / 11Tutorial 10Thursday, 16 December 2010, 13.45–15.15, T 212

Let V, H be separable Hilbert spaces. We consider the abstract problem: find  $u \in H^1((0,T),V;H)$  such that

$$\frac{d}{dt}\underbrace{(u(t), v)_H}_{=\langle u'(t), v \rangle} + a(u(t), v) = \langle F(t), v \rangle \qquad \forall v \in V \quad \forall t \in (0, T) \text{ a.e.,} \\
\underbrace{u(0)}_{=\langle u'(t), v \rangle} u(0) = u_0 \qquad \text{in } H,$$
(10.1)

for given  $u_0 \in H$  and  $F \in L^2((0,T), V^*)$ , where  $(V, H, V^*)$  is an evolution triple, i.e.,

- $V \subset H$ ,
- there exists a constant c > 0 such that  $||v||_H \le c ||v||_V$  for all  $v \in V$ ,
- V is dense in H.

Recall that the weak time derivative  $u' \in L^2((0, T), V^*)$  fulfills

$$\int_0^T \varphi(t) \langle u'(t), v \rangle dt = -\int_0^T \varphi'(t) (u(t), v)_H dt \qquad \forall v \in V \ \forall \varphi \in C_0^\infty(0, T).$$

56 Show that for all  $\lambda \in \mathbb{R}$ : the function  $u \in H^1((0,T), V; H)$  is a solution of (10.1) if and only if  $u_{\lambda} \in H^1((0,T), V; H)$  solves

$$\frac{d}{dt}(u_{\lambda}(t), v)_{H} + a_{\lambda}(u_{\lambda}(t), v) = \langle F_{\lambda}(t), v \rangle \quad \forall v \in V \quad \forall t \in (0, T) \text{ a.e.,} \\
u_{\lambda}(0) = u_{0} \qquad \text{in } H,$$
(10.2)

where

$$u_{\lambda}(t) = e^{-\lambda t} u(t), \qquad a_{\lambda}(w, v) = a(w, v) + \lambda(w, v)_H, \qquad F_{\lambda}(t) = e^{-\lambda t} F(t).$$

*Hint*: use the definition of the weak time derivative u'(t) to compute  $u'_{\lambda}(t)$ .

57 Let  $a: V \times V \to \mathbb{R}$  be a bounded bilinear form and let there exist constants  $\lambda \in \mathbb{R}$ and  $\mu_1 > 0$  such that the so-called *Gårding inequality*)

$$a(v,v) + \lambda ||v||_{H}^{2} \ge \mu_{1} ||v||_{V}^{2} \quad \forall v \in V$$

holds. Show that problem (10.1) is uniquely solvable for any  $F \in L^2((0,T); V^*)$  and  $u_0 \in H$ .

*Hint:* Use Exercise  $\lfloor 56 \rfloor$  and Theorem 2.8.

58 Consider the bilinear form

$$a(w, v) := \int_0^1 a(x) \frac{\partial w}{\partial x}(x) \frac{\partial v}{\partial x}(x) + b(x) \frac{\partial w}{\partial x}(x) v(x) + c(x) w(x) v(x) dx$$

in  $H^1(0, 1)$  with  $a, b, c \in L^{\infty}(0, 1)$ , where  $a_0 := \inf_{x \in (0, 1)} a(x) > 0$ . Show the Gårding inequality: there exist constants  $\lambda \in \mathbb{R}$  and  $\mu_1 > 0$  such that

$$a(v, v) + \lambda \|v\|_{L^2(0,1)}^2 \ge \mu_1 \|v\|_{H^1(0,1)}^2 \quad \forall v \in H^1(0,1).$$

*Hint:* Choose  $\lambda$  such that the assumptions of Excercise [11] (Tutorial 3) hold for the bilinear form  $a_{\lambda}(w, v) := a(w, v) + \lambda(w, v)_{L^2(0,1)}$ .

59 Let  $C^1([0,T], V)$  denote the space of continuous functions in [0,T] with values in the Hilbert space V that have a continuous *classical* derivative, i. e., for  $v \in C^1([0,T], V)$  the limit

$$v'(t) := \lim_{\tau \to 0} \frac{1}{\tau} (v(t+\tau) - v(t))$$

exists for all  $t \in [0, T]$  and the function  $v' : [0, T] \to V$  is continuus. Show that for all  $s, t \in [0, T]$  and for all  $v \in C^1([0, T], V)$ :

$$\frac{1}{2} \left( v(t), v(t) \right)_{H} = \frac{1}{2} \left( v(s), v(s) \right)_{H} + \int_{s}^{t} \left( v'(\sigma), v(\sigma) \right)_{H} d\sigma.$$
(10.3)

*Hint:* Prove and use the identity

$$\frac{1}{2}\left[\left(v(\sigma), v(\sigma)\right)_{H}\right]' = \left(v'(\sigma), v(\sigma)\right)_{H}.$$

60 Prove the statement of Lemma 2.7 for continuously differentiable functions, i.e., show that there exists a constant C > 0 such that

$$\max_{t \in [0,T]} \|v(t)\|_{H} \leq C \|v\|_{H^{1}((0,T),V;H)} \qquad \forall v \in C^{1}([0,T],V).$$

*Hint:* Integrate identity (10.3) w.r.t. s over [0, T]. Note that  $||v'||^2_{L^2((0,T),V^*)}$  is the integral over the (square of the)  $V^*$ -norm of the mapping  $w \mapsto (v'(t), w)_H$ . Show and use that

$$(v'(\sigma), w)_H \leq \|v'(\sigma)\|_{V^*} \|w\|_V \leq \frac{1}{2} \left[ \|v'(\sigma)\|_{V^*}^2 + \|w\|_{V^*}^2 \right] \quad \forall w \in V.$$

61 Consider the Courant FE semi-discretization of our 1D parabolic model problem  $\frac{\partial u}{\partial t}(x,t) - \frac{\partial^2 u}{\partial x^2}(x,t) = f(x,t)$  for all  $(x,t) \in (0,1) \times (0,T)$ , with u(0,t) = 0 and  $\frac{\partial u}{\partial x}(1,t) = g_1(t)$  for all  $t \in [0,T]$ .

Derive an upper bound for  $||M_h^{-1}K_h||$  where  $||\cdot||$  is the operator norm w.r.t.  $||\cdot||_{\ell^2}$ . This will imply that the right hand side of problem (5) from the lecture satisfies the assumptions of the Picard-Lindelöf Theorem, and so (5) is uniquely solvable.

*Hint:*  $M_h^{-1}K_h$  is positive definite with respect to the inner product  $(\underline{v}_h, \underline{w}_h)_{M_h} := (M_h \underline{v}_h, \underline{w}_h)_{\ell^2}$ . Use the Rayleigh quotient and the eigenvalue bounds from Chapter 1 to get an upper bound for  $\lambda_{\max}(M_h^{-1}K_h)$ .