

Towards a IETI-DP Solver on Non-Matching Multi-Patch Domains

Rainer Schneckleitner

Institute of Computational Mathematics, Johannes Kepler University
Altenberger Str. 69, 4040 Linz, Austria

Stefan Takacs

Johann Radon Institute for Computational and Applied Mathematics
Austrian Academy of Sciences
Altenberger Str. 69, 4040 Linz, Austria

NuMa-Report No. 2021-02

March 2021

Technical Reports before 1998:

1995

- 95-1 Hedwig Brandstetter
Was ist neu in Fortran 90? March 1995
- 95-2 G. Haase, B. Heise, M. Kuhn, U. Langer
Adaptive Domain Decomposition Methods for Finite and Boundary Element Equations. August 1995
- 95-3 Joachim Schöberl
An Automatic Mesh Generator Using Geometric Rules for Two and Three Space Dimensions. August 1995

1996

- 96-1 Ferdinand Kickingger
Automatic Mesh Generation for 3D Objects. February 1996
- 96-2 Mario Goppold, Gundolf Haase, Bodo Heise und Michael Kuhn
Preprocessing in BE/FE Domain Decomposition Methods. February 1996
- 96-3 Bodo Heise
A Mixed Variational Formulation for 3D Magnetostatics and its Finite Element Discretisation. February 1996
- 96-4 Bodo Heise und Michael Jung
Robust Parallel Newton-Multilevel Methods. February 1996
- 96-5 Ferdinand Kickingger
Algebraic Multigrid for Discrete Elliptic Second Order Problems. February 1996
- 96-6 Bodo Heise
A Mixed Variational Formulation for 3D Magnetostatics and its Finite Element Discretisation. May 1996
- 96-7 Michael Kuhn
Benchmarking for Boundary Element Methods. June 1996

1997

- 97-1 Bodo Heise, Michael Kuhn and Ulrich Langer
A Mixed Variational Formulation for 3D Magnetostatics in the Space $H(\text{rot}) \cap H(\text{div})$ February 1997
- 97-2 Joachim Schöberl
Robust Multigrid Preconditioning for Parameter Dependent Problems I: The Stokes-type Case. June 1997
- 97-3 Ferdinand Kickingger, Sergei V. Nepomnyaschikh, Ralf Pfau, Joachim Schöberl
Numerical Estimates of Inequalities in $H^{\frac{1}{2}}$. August 1997
- 97-4 Joachim Schöberl
Programmbeschreibung NAOMI 2D und Algebraic Multigrid. September 1997

From 1998 to 2008 technical reports were published by SFB013. Please see

<http://www.sfb013.uni-linz.ac.at/index.php?id=reports>

From 2004 on reports were also published by RICAM. Please see

<http://www.ricam.oeaw.ac.at/publications/list/>

For a complete list of NuMa reports see

<http://www.numa.uni-linz.ac.at/Publications/List/>

Towards a IETI-DP solver on non-matching multi-patch domains

Rainer Schneckleitner* Stefan Takacs

Abstract Recently, the authors have proposed and analyzed isogeometric tearing and interconnecting (IETI-DP) solvers for multi-patch discretizations in Isogeometric Analysis. Conforming and discontinuous Galerkin settings have been considered. In both cases, we have assumed that the interfaces between the patches consist of whole edges. In this paper, we present a generalization that allows us to drop this requirement. This means that the patches can meet in T-junctions, which increases the flexibility of the geometric model significantly. We use vertex-based primal degrees of freedom. For the T-junctions, we propose to follow the idea of “fat vertices”.

1 Introduction

Isogeometric Analysis (IgA), see [6], is a method for discretizing partial differential equations (PDEs). The goal of its development has been to enhance the interface between computer-aided design (CAD) and simulation. Current state-of-the-art CAD tools use B-splines and NURBS for the representation of the computational domain. In IgA, the same kind of bases is also utilized to discretize the PDEs. Complex domains for real-world applications are usually the union of many patches, parametrized with individual geometry functions (multi-patch IgA). We focus on non-overlapping patches.

Rainer Schneckleitner

Institute of Computational Mathematics, Johannes Kepler University Linz, Altenberger Straße 69, 4040 Linz, Austria e-mail: schneckleitner@numa.uni-linz.ac.at,
Corresponding author

Stefan Takacs

RICAM, Austrian Academy of Sciences, Altenberger Straße 69, 4040 Linz, Austria e-mail: stefan.takacs@ricam.oeaw.ac.at

If the grids are not conforming and/or the interfaces between the patches do not consist of whole edges then discontinuous Galerkin (dG) methods are the discretization techniques of choice. A well studied representative is the symmetric interior discontinuous Galerkin (SIPG) method, cf. [1]. It has already been adapted and analyzed in IgA, cf. [8, 9, 14] and others. An obvious choice to solve discretized PDEs on domains with many non-overlapping patches are tearing and interconnecting methods. The variant we are interested in is the dual-primal approach, see [3] for FETI-DP and [7, 4, 5] for its extension to IgA, which is called accordingly dual-primal isogeometric tearing and interconnecting method (IETI-DP). In [13, 14], the authors have presented a p - and h -robust convergence analysis. The authors have assumed that the interfaces consist of whole edges. If the vertices are chosen as primal degrees of freedom, it was shown that the condition number of the preconditioned Schur complement system is, under proper assumptions, bounded by

$$C p \left(1 + \log p + \max_{k=1, \dots, K} \log \frac{H_k}{h_k} \right)^2, \quad (1)$$

where p is the spline degree, h_k is the grid size on patch $\Omega^{(k)}$ and H_k is the diameter of $\Omega^{(k)}$ and $C > 0$ is a constant independent of these quantities. In this paper, we construct a new IETI-DP method that can deal with interfaces that do not consist of whole edges. This means that the patches can meet in T-junctions, which increases the flexibility of the geometric model significantly. In this IETI-DP variant, the construction of the coarse space is based on the idea of “fat vertices”: We consider every basis function that is supported on a vertex or T-junction as primal degree of freedom. The numerical experiments indicate that a similar condition number bound to (1) might hold.

The remainder of this paper is organized as follows. In Section 2 we describe the model problem. In Section 3 we introduce the IETI-DP solver and we end this paper with numerical experiments in Section 4.

2 The problem setting

Let $\Omega \subset \mathbb{R}^2$ be open, simply connected and bounded with Lipschitz boundary $\partial\Omega$. $L_2(\Omega)$ and $H^1(\Omega)$ are the common Lebesgue and Sobolev spaces. As usual, $H_0^1(\Omega) \subset H^1(\Omega)$ denotes the subspace of functions that vanish on $\partial\Omega$.

We consider the following model problem: Find $u \in H_0^1(\Omega)$ such that

$$\int_{\Omega} \nabla u \cdot \nabla v \, dx = \int_{\Omega} f v \, dx \quad \text{for all } v \in H_0^1(\Omega) \quad (2)$$

with a given source function $f \in L_2(\Omega)$. We assume that Ω is a composition of K non-overlapping patches $\Omega^{(k)}$, where every patch $\Omega^{(k)}$ is parametrized by a geometry function

$$G_k : \widehat{\Omega} := (0, 1)^2 \rightarrow \Omega^{(k)} := G_k(\widehat{\Omega}) \subset \mathbb{R}^2, \quad (3)$$

that has a continuous extension to the closure of $\widehat{\Omega}$ and such that $\nabla G_k \in L_\infty(\widehat{\Omega})$ and $(\nabla G_k)^{-1} \in L_\infty(\widehat{\Omega})$.

We consider the case where the pre-images of the (Dirichlet) boundary consist of whole edges. The indices of neighboring patches $\Omega^{(\ell)}$ of $\Omega^{(k)}$, that share at least a part of their boundaries, is collected in the set

$$\mathcal{N}_\Gamma(k) := \{\ell \neq k : \text{meas}(\partial\Omega^{(k)} \cap \partial\Omega^{(\ell)}) > 0\},$$

where $\text{meas } T$ is the measure of T . For any $\ell \in \mathcal{N}_\Gamma(k)$, we write $\Gamma^{(k,\ell)} = \partial\Omega^{(k)} \cap \partial\Omega^{(\ell)}$. The endpoints of $\partial\Omega^{(k)} \cap \partial\Omega^{(\ell)}$ that are not located on the (Dirichlet) boundary of Ω are referred to as junctions. A junction could be a common vertex or a T-junction.

For the IgA discretization spaces, we first construct a B-spline space $\widehat{V}^{(k)}$ on the parameter domain $\widehat{\Omega}$ by tensorization of two univariate B-spline spaces. The function spaces on the physical domain are then defined by the pull-back principle: $V^{(k)} := \widehat{V}^{(k)} \circ G_k^{-1}$.

The product of the local spaces gives the global approximation space $V := V^{(1)} \times \dots \times V^{(K)}$. On this discretization space, we could introduce the SIPG formulation, cf. [1, 14]. Since we are interested in a domain decomposition approach, we need patch-local formulations of SIPG.

3 The dG IETI-DP solver

For those patch-local formulations, we adapt the ideas of [2, 5, 4] and others. We choose local function spaces $V_e^{(k)}$ to be the product space of $V^{(k)}$ and the neighboring trace spaces $V^{(k,\ell)}$, which are the restrictions of $V^{(\ell)}$ to $\Gamma^{(k,\ell)}$. A function $v_e^{(k)} \in V_e^{(k)}$ is represented as a tuple $v_e^{(k)} = (v^{(k)}, (v^{(k,\ell)})_{\ell \in \mathcal{N}_\Gamma(k)})$, where $v^{(k)} \in V^{(k)}$ and $v^{(k,\ell)} \in V^{(k,\ell)}$. Note that the traces of the basis functions for $V^{(\ell)}$ restricted to $\Gamma^{(k,\ell)}$ form a basis of $V^{(k,\ell)}$. The basis for $V_e^{(k)}$ consists of the basis functions of $V^{(k)}$ and the basis functions for $V^{(k,\ell)}$. The basis functions on $V^{(k,\ell)}$ are usually visualized as living on artificial interfaces.

On each patch, we consider the local problem: Find $u_e^{(k)} \in V_e^{(k)}$ such that

$$a_e^{(k)}(u_e^{(k)}, v_e^{(k)}) = \langle f_e^{(k)}, v_e^{(k)} \rangle \quad \text{for all } v_e^{(k)} \in V_e^{(k)},$$

where

$$\begin{aligned} a_e^{(k)}(u_e^{(k)}, v_e^{(k)}) &:= a^{(k)}(u_e^{(k)}, v_e^{(k)}) + m^{(k)}(u_e^{(k)}, v_e^{(k)}) + r^{(k)}(u_e^{(k)}, v_e^{(k)}), \\ \langle f_e^{(k)}, v_e^{(k)} \rangle &:= \int_{\Omega^{(k)}} f v^{(k)} dx, \\ a^{(k)}(u_e^{(k)}, v_e^{(k)}) &:= \int_{\Omega^{(k)}} \nabla u^{(k)} \cdot \nabla v^{(k)} dx, \\ m^{(k)}(u_e^{(k)}, v_e^{(k)}) &:= \sum_{\ell \in \mathcal{N}_\Gamma^{(k)}} \int_{\Gamma^{(k, \ell)}} \frac{\partial u^{(k)}}{\partial n_k} (v^{(k, \ell)} - v^{(k)}) ds, \\ &\quad + \sum_{\ell \in \mathcal{N}_\Gamma^{(k)}} \int_{\Gamma^{(k, \ell)}} \frac{\partial v^{(k)}}{\partial n_k} (u^{(k, \ell)} - u^{(k)}) ds, \\ r^{(k)}(u_e^{(k)}, v_e^{(k)}) &:= \sum_{\ell \in \mathcal{N}_\Gamma^{(k)}} \int_{\Gamma^{(k, \ell)}} \frac{\delta p^2}{h_{k\ell}} (u^{(k, \ell)} - u^{(k)}) (v^{(k, \ell)} - v^{(k)}) ds \end{aligned}$$

and n_k denotes the outward unit normal vector and δ is the dG penalty parameter, which has to be chosen large enough in order to guarantee that the bilinear form $a_e^{(k)}(\cdot, \cdot)$ is coercive. In [15], it was shown that δ can be chosen independently of p .

The discretization of $a_e^{(k)}(\cdot, \cdot)$ and $\langle f_e^{(k)}, \cdot \rangle$ gives a local system, which we write as

$$\begin{pmatrix} A_{\text{II}}^{(k)} & A_{\text{I}\Gamma}^{(k)} \\ A_{\Gamma\text{I}}^{(k)} & A_{\Gamma\Gamma}^{(k)} \end{pmatrix} \begin{pmatrix} \underline{u}_{\text{I}}^{(k)} \\ \underline{u}_{\Gamma}^{(k)} \end{pmatrix} = \begin{pmatrix} \underline{f}_{\text{I}}^{(k)} \\ \underline{f}_{\Gamma}^{(k)} \end{pmatrix}, \quad (4)$$

where the index I refers to the basis functions that are only supported in the interior of $\Omega^{(k)}$ and the index Γ refers to the remaining basis functions, i.e., those living on the patch boundary and on the artificial interfaces. We eliminate the interior degrees of freedom in (4) for every $k = 1, \dots, K$ to get the block diagonal Schur complement system

$$S \underline{w} = \underline{g}, \quad (5)$$


where the individual blocks of S are given by $S^{(k)} = A_{\Gamma\Gamma}^{(k)} - A_{\Gamma\text{I}}^{(k)} (A_{\text{II}}^{(k)})^{-1} A_{\text{I}\Gamma}^{(k)}$.

The IETI-DP method requires carefully selected primal degrees of freedom to be solvable. We choose the degrees of freedom associated to the basis functions which are non-zero on a junction to be primal. For every standard corner, we only have one primal degree of freedom per patch, as in [14]. On a T-junction however, the number of non-zero basis functions grows linearly with p . Since we take all of them, we refer to ‘‘fat vertices’’ in this context.

$C = \text{diag}(C^{(1)}, \dots, C^{(K)})$ is the constraint matrix, i.e., it is defined such that $Cw = 0$ if and only if the associated function w vanishes at the primal degrees of freedom. The matrix Ψ represents the energy minimizing basis functions for the space of primal degrees of freedom.

Furthermore, we introduce the jump matrix B , which models the jumps of the functions between the patch boundaries and the associated artificial interfaces. Each row corresponds to one degree of freedom (coefficient for a basis function) on the patch boundary and one artificial interface; as usual, each row has only two non-zero coefficients that are -1 and 1 . Primal degrees of freedom are excluded. For a visualization, see Fig. 1, where the primal degrees of freedom are marked with solid lines and the dotted arrows show the action of the jump matrix B . The basis functions on the artificial interfaces are labeled with the same symbols from the original spaces.

Fig. 1: Action of matrix B (dotted lines) and primal degrees of freedom (solid lines)



The following problem is equivalent to the SIPG discretization of (2), cf. [10]: Find $(\underline{w}_\Delta, \underline{\mu}, \underline{w}_\Pi, \underline{\lambda})$ such that

$$\begin{pmatrix} S & C^\top & & B^\top \\ C & & & \\ & \Psi^\top S \Psi & & (B\Psi)^\top \\ B & & B\Psi & \end{pmatrix} \begin{pmatrix} \underline{w}_\Delta \\ \underline{\mu} \\ \underline{w}_\Pi \\ \underline{\lambda} \end{pmatrix} = \begin{pmatrix} \underline{g} \\ 0 \\ \Psi^\top \underline{g} \\ 0 \end{pmatrix}.$$

We obtain the solution of the original problem by $\underline{w} = \underline{w}_\Delta + \Psi \underline{w}_\Pi$. We build a Schur complement of this system to get the linear problem

$$F \underline{\lambda} = \underline{d}. \quad (6)$$

We solve (6) with a preconditioned conjugate gradient (PCG) solver with the scaled Dirichlet preconditioner

$$M_{\text{sD}} := BD^{-1}SD^{-1}B^\top,$$

where D is a diagonal matrix defined based on the principle of multiplicity scaling, cf. [13, 12].

4 Numerical results

We consider the model problem

$$\begin{aligned} -\Delta u(x, y) &= 2\pi^2 \sin(\pi x) \sin(\pi y) && \text{for } (x, y) \in \Omega \\ u &= 0 && \text{on } \partial\Omega, \end{aligned}$$

on the geometries depicted in Fig. 2. Both represent the same computational domain with an inner radius of 1 and an outer radius of 2. The ring in Fig. 2a consists of 20 patches each of which has a width of 0.2. For the ring in Fig. 2b, the thin layer of patches has a width of 0.02, the other layers have a correspondingly larger width. We use NURBS of degree 2 to parametrize all patches. In the coarsest setting, i.e., $r = 0$, the discretization spaces on all patches consist of global polynomials only. The discretization spaces for $r = 1, 2, 3, \dots$ are obtained by uniform refinement steps. We use a PCG solver to solve system (6) with the preconditioner M_{sD} and to estimate the condition number $\kappa(M_{\text{sD}}F)$, where we use the zero vector as initial guess. All experiments are carried out in the C++ library G+Smo, cf. [11] and are executed on the Radon1¹ cluster in Linz.




Fig. 2: Computational domains and the decomposition into patches

In the Table 1, we report on the iteration counts (it) and the condition numbers (κ) for various refinement levels r and various spline degrees p , where we chose C^s -smoothness with $s = p - 1$ within the patches. The tables show the expected behavior with respect to h . The condition number decreases when we increase the spline degree p , which is better than one would expect from the theory in [14]. Although the width of the thin patches in Fig. 2b is one tenth of the width of the patches in Fig. 2a, the condition number grows only by a factor between 5 and 6. Also the iteration counts grow only mildly.

The Table 2 presents the parallel solving times for n processors. We only consider the domain in Fig. 2a again with $s = p - 1$. We see that the speedup

¹ <https://www.ricam.oeaw.ac.at/hpc/>

r	Fig. 2a								Fig. 2b							
	$p = 2$		$p = 3$		$p = 6$		$p = 7$		$p = 2$		$p = 3$		$p = 6$		$p = 7$	
	it	κ	it	κ	it	κ	it	κ	it	κ	it	κ	it	κ	it	κ
4	9	3.7	9	3.5	8	2.4	8	2.1	12	18.0	13	18.0	13	12.9	11	11.6
5	10	4.6	10	4.5	9	3.8	9	3.5	20	24.1	19	23.4	19	19.3	18	18.0
6	10	5.8	10	5.5	10	4.9	10	4.8	22	31.7	22	29.9	21	25.9	20	24.8
7	11	6.3	11	6.2	10	5.6	10	5.5	24	37.2	24	36.3	22	31.3	22	30.1
8	11	6.7	11	6.7	11	6.3	10	5.6	24	43.2	24	42.3	24	36.5	24	31.6

Table 1: Iterations (it) and condition numbers (κ)

rate with respect to n is a bit smaller than the expected rate of 2. This is probably caused by the rather small number of patches in the computational domain.

r	$p = 3$					$p = 7$				
	$n = 1$	$n = 2$	$n = 4$	$n = 8$	$n = 16$	$n = 1$	$n = 2$	$n = 4$	$n = 8$	$n = 16$
6	3.8	2.8	2.4	1.2	0.8	10.0	6.5	5.0	2.5	1.75
7	24.0	16.1	13.6	6.4	4.1	47.0	31.7	26.1	12.2	9.3
8	107.0	81.4	66.8	29.5	19.5	220.0	158.7	129.4	56.7	45.4

Table 2: Solving times (sec.); Fig. 2a

In Table 3 we report on the iteration counts and the condition numbers for the decomposition in Fig. 2a when we change the smoothness s of the B-splines within the patches. The numbers in the table show the behavior for $r = 5$. We see that for a fixed smoothness s the condition number grows slightly with respect to the spline degree p . For a fixed degree p , we observe a decline in the condition number when we increase the smoothness s .

Acknowledgements The first author was supported by the Austrian Science Fund (FWF): S117 and W1214-04. Also, the second author has received support from the Austrian Science Fund (FWF): P31048.

References

1. D. Arnold. An interior penalty finite element method with discontinuous elements. *SIAM J. Numer. Anal.*, 19(4):742 – 760, 1982.

s	p = 2		p = 3		p = 4		p = 5		p = 6		p = 7	
	it	κ	it	κ	it	κ	it	κ	it	κ	it	κ
0	10	5.0	10	5.3	10	5.4	10	5.5	10	5.6	10	5.6
1	10	4.6	10	5.2	10	5.3	10	5.4	10	5.5	10	5.5
2			10	4.5	10	5.0	10	5.3	10	5.4	10	5.5
3					9	4.2	10	4.9	10	5.1	10	5.3
4							9	4.0	10	4.7	10	5.0
5									9	3.8	9	4.5
6											9	3.5

Table 3: Iterations (it) and condition number (κ); $r = 5$; Fig. 2a

2. M. Dryja, J. Galvis, and M. Sarkis. A FETI-DP Preconditioner for a Composite Finite Element and Discontinuous Galerkin Method. *SIAM J. Numer. Anal.*, 51(1):400–422, 2013.
3. C. Farhat, M. Lesoinne, P. L. Tallec, K. Pierson, and D. Rixen. FETI-DP: A dual-primal unified FETI method I: A faster alternative to the two-level FETI method. *Int. J. Numer. Methods Eng.*, 50:1523–1544, 2001.
4. C. Hofer. Analysis of discontinuous Galerkin dual-primal isogeometric tearing and interconnecting methods. *Math. Models Methods Appl. Sci.*, 28(1):131–158, 2018.
5. C. Hofer and U. Langer. Dual-primal isogeometric tearing and interconnecting solvers for multipatch continuous and discontinuous Galerkin IgA equations. *PAMM*, 16(1):747–748, 2016.
6. T. J. R. Hughes, J. A. Cottrell, and Y. Bazilevs. Isogeometric analysis: CAD, finite elements, NURBS, exact geometry and mesh refinement. *Comput. Methods Appl. Mech. Eng.*, 194(39-41):4135 – 4195, 2005.
7. S. Kleiss, C. Pechstein, B. Jüttler, and S. Tomar. IETI-Isogeometric Tearing and Interconnecting. *Comput. Methods Appl. Mech. Eng.*, 247-248:201–215, 2012.
8. U. Langer, A. Mantzaflaris, S. E. Moore, and I. Touloupoulos. Multipatch Discontinuous Galerkin Isogeometric Analysis. In B. Jüttler and B. Simeon, editors, *Isogeometric Analysis and Applications 2014*, pages 1–32. Springer International Publishing, 2015.
9. U. Langer and I. Touloupoulos. Analysis of multipatch discontinuous Galerkin IgA approximations to elliptic boundary value problems. *Comp. Vis. Sci.*, 17(5):217 – 233, 2015.
10. J. Mandel, C. R. Dohrmann, and R. Tezaur. An algebraic theory for primal and dual substructuring methods by constraints. *Appl. Numer. Math.*, 54(2):167–193, 2005.
11. A. Mantzaflaris, R. Schneckleitner, S. Takacs, and others (see website). G+Smo (Geometry plus Simulation modules). <http://github.com/gismo>, 2020.
12. C. Pechstein. *Finite and Boundary Element Tearing and Interconnecting Solvers for Multiscale Problems*. Springer, Heidelberg, 2013.
13. R. Schneckleitner and S. Takacs. Condition number bounds for IETI-DP methods that are explicit in h and p . *Math. Models Methods Appl. Sci.*, 30(11):2067 – 2103, 2020.
14. R. Schneckleitner and S. Takacs. Convergence theory for IETI-DP solvers for discontinuous Galerkin Isogeometric Analysis that is explicit in h and p , 2020. Submitted. <https://arxiv.org/pdf/2005.09546.pdf>.
15. S. Takacs. A quasi-robust discretization error estimate for discontinuous Galerkin Isogeometric Analysis, 2019. Submitted. <https://arxiv.org/pdf/1901.03263.pdf>.

Latest Reports in this series

2009 - 2018

[..]

2019

- | | | |
|---------|---|---------------|
| 2019-01 | Helmut Gfrerer and Jiří V. Outrata
<i>On a Semismooth* Newton Method for Solving Generalized Equations</i> | April 2019 |
| 2019-02 | Matúš Benko, Michal Červinka and Tim Hoheisel
<i>New Verifiable Sufficient Conditions for Metric Subregularity of Constraint Systems with Applications to Disjunctive Programs</i> | June 2019 |
| 2019-03 | Matúš Benko
<i>On Inner Calmness*, Generalized Calculus, and Derivatives of the Normal-cone Map</i> | October 2019 |
| 2019-04 | Rainer Schnockenleitner and Stefan Takacs
<i>Condition number bounds for IETI-DP methods that are explicit in h and p</i> | December 2019 |
| 2019-05 | Clemens Hofreither, Ludwig Mitter and Hendrik Speleers
<i>Local multigrid solvers for adaptive Isogeometric Analysis in hierarchical spline spaces</i> | December 2019 |

2020

- | | | |
|---------|---|---------------|
| 2020-01 | Ioannis Touloupoulos
<i>Viscoplastic Models and Finite Element Schemes for the Hot Rolling Metal Process</i> | February 2020 |
| 2020-02 | Rainer Schnockenleitner and Stefan Takacs
<i>Convergence Theory for IETI-DP Solvers for Discontinuous Galerkin Isogeometric Analysis That Is Explicit in h and p</i> | May 2020 |
| 2020-03 | Svetoslav Nakov and Ioannis Touloupoulos
<i>Convergence Estimates of Finite Elements for a Class of Quasilinear Elliptic Problems</i> | May 2020 |
| 2020-04 | Helmut Gfrerer, Jiří V. Outrata and Jan Valdman
<i>On the Application of the Semismooth* Newton Method to Variational Inequalities of the Second Kind</i> | July 2020 |

2021

- | | | |
|---------|--|---------------|
| 2020-01 | Ioannis Touloupoulos
<i>A Continuous Space-Time Finite Element Scheme for Quasilinear Parabolic Problems</i> | February 2021 |
| 2020-02 | Rainer Schnockenleitner and Stefan Takacs
<i>Towards a IETI-DP Solver on Non-Matching Multi-Patch Domains</i> | March 2021 |

From 1998 to 2008 reports were published by SFB013. Please see

<http://www.sfb013.uni-linz.ac.at/index.php?id=reports>

From 2004 on reports were also published by RICAM. Please see

<http://www.ricam.oeaw.ac.at/publications/>

For a complete list of NuMa reports see

<http://www.numa.uni-linz.ac.at/Publications/List/>